

# A note on bicomplex Orlicz spaces

By

S. Dubey, R. Kumar and K. Sharma

## Abstract

In this paper we study bicomplex function spaces and in particular we study bicomplex Orlicz spaces.

## 1 INTRODUCTION

In this section we summarize some basic properties already established for the bicomplex numbers and also give some remarks and conclusions. The set  $\mathbb{BC}$  of bicomplex numbers is defined as

$$\mathbb{BC} = \{z_1 + z_2j : z_1, z_2 \in \mathbb{C}(i)\},$$

where  $i$  and  $j$  are independent imaginary units such that  $i^2 = j^2 = -1$  and  $\mathbb{C}(i) = \{x + iy : x, y \in \mathbb{R}\}$ . The product of any two units is commutative and satisfies

$$ij = ji = k, ik = -j, jk = -i.$$

The set of bicomplex numbers,  $\mathbb{BC}$ , can also be defined as

$$\mathbb{BC} = \{a + bi + cj + dk : a, b, c, d, \in \mathbb{R}\}.$$

Under addition and multiplication defined as :

$$\begin{aligned} Z + W &= (z_1 + z_2j) + (w_1 + w_2j) \\ &= (z_1 + w_1) + (z_2 + w_2)j \\ Z.W &= (z_1 + z_2j).(w_1 + w_2j) \\ &= (z_1w_1 - z_2w_2) + (z_1w_2 + z_2w_1)j \end{aligned}$$

$\mathbb{BC}$  becomes a commutative ring with unity and therefore a module over itself. Also since the field  $\mathbb{C}(i)$  is a subring of  $\mathbb{BC}$ , therefore  $\mathbb{BC}$  can also be seen as a vector space over  $\mathbb{C}(i)$ . If we put  $z_1 = x$  and  $z_2 = iy$  with  $x, y \in \mathbb{R}$ , then we obtain the set of hyperbolic numbers

$$\mathbb{D} = \{x + yk : k^2 = 1, x, y \in \mathbb{R}\}.$$

Three types of conjugations can be defined on the set of bicomplex numbers. With  $Z = z_1 + z_2j$ , we define

- (i)  $\overline{Z} = \overline{z}_1 + \overline{z}_2 j$
- (ii)  $Z^\dagger = z_1 - z_2 j$
- (iii)  $Z^* = (\overline{Z})^* = \overline{z}_1 - \overline{z}_2 j$ .

Each conjugate is additive, multiplicative and involute on  $\mathbb{BC}$ . Let  $e = \frac{1 + ij}{2}$ . Then the  $\dagger$ -conjugate of  $e$  is given by  $e^\dagger = \frac{1 - ij}{2}$ . Any bicomplex number  $Z = z_1 + z_2 j$  can be uniquely expressed as  $Z = \beta_1 e + \beta_2 e^\dagger$ , where  $\beta_1 = z_1 - z_2 i$  and  $\beta_2 = z_1 + z_2 i$  are in  $\mathbb{C}(i)$ . The bicomplex numbers  $e$  and  $e^\dagger$  are hyperbolic numbers with

$$\begin{aligned} e^2 &= e; & e^* &= e; & e + e^\dagger &= 1; \\ (e^\dagger)^2 &= e^\dagger; & (e^\dagger)^* &= e^\dagger; & e \cdot e^\dagger &= 0. \end{aligned}$$

$e$  and  $e^\dagger$  form the idempotent basis of bicomplex numbers. The uniqueness of idempotent representation of bicomplex numbers allows the introduction of two projection operators  $\pi_{1,i} : \mathbb{BC} \mapsto \mathbb{C}(i)$  and  $\pi_{2,i} : \mathbb{BC} \mapsto \mathbb{C}(i)$  defined as

$$\pi_{l,i}(Z) = \pi_{l,i}(\beta_1 e + \beta_2 e^\dagger) = \beta_l \in \mathbb{C}(i) \text{ for } l = 1, 2,$$

with the property that  $\pi_{1,i}e + \pi_{2,i}e^\dagger = I$  and both the projection maps are additive and multiplicative, i.e.  $\pi_{l,i}(Z + W) = \pi_{l,i}(Z) + \pi_{l,i}(W)$  and  $\pi_{l,i}(Z.W) = \pi_{l,i}(Z) \cdot \pi_{l,i}(W)$  for all  $Z, W \in \mathbb{BC}$  and  $l = 1, 2$ .

For  $Z = z_1 e + z_2 e^\dagger \in \mathbb{BC}$ , the norm of  $Z$  is defined as  $\|Z\| = \frac{1}{\sqrt{2}} \sqrt{|z_1|^2 + |z_2|^2}$ .

Let  $A \subseteq \mathbb{BC}$ . Also each  $x \in A$  is of the form  $Z = z_1 e + z_2 e^\dagger$  where  $z_1$  and  $z_2$  are in  $\mathbb{C}(i)$ . Corresponding to each  $A \subseteq \mathbb{BC}$ , we associate two subsets of  $\mathbb{C}(i)$ , viz.  $A_e$  and  $A_{e^\dagger}$  comprising of  $\pi_{1,i}(Z)$  and  $\pi_{2,i}(Z)$ , respectively, for all  $Z \in A$ . Therefore  $A_e = \pi_{1,i}(A) = eA$  and  $A_{e^\dagger} = \pi_{2,i}(A) = e^\dagger A$ .

Also let  $A_1$  and  $A_2$  be two sets defined as

$$\begin{aligned} A_1 &= \{z_1 - iz_2 : z_1, z_2 \in \mathbb{C}(i)\} \\ A_2 &= \{z_1 + iz_2 : z_1, z_2 \in \mathbb{C}(i)\}. \end{aligned}$$

Then  $A_1$  and  $A_2$  are copies of  $\mathbb{C}(i)$  such that  $\mathbb{BC} = A_1 e + A_2 e^\dagger$ . Therefore for any subset  $U$  of  $\mathbb{BC}$ , we have  $U = U_1 e + U_2 e^\dagger$ , where  $U_1$  and  $U_2$  are subsets of  $A_1$  and  $A_2$  respectively. It is easy to show now that the characteristic functions  $\chi_U$  is given by  $\chi_U = \chi_{U_1} \chi_{U_2} e + \chi_{U_1} \chi_{U_2} e^\dagger$ .

**Remark 1.1.**  $\mathbb{BC}$  together with the norm defined above form a generalized normed algebra, since  $\|Z.W\| \leq \frac{1}{\sqrt{2}} \|Z\| \cdot \|W\|$ . Also since  $\mathbb{BC} \simeq \mathbb{R}^4$  and  $\mathbb{R}^4$  is complete with respect to usual metric, it follows that  $\mathbb{BC}$  forms a generalized Banach algebra. The product of two bicomplex numbers  $Z = \beta_{Z,1} e + \beta_{Z,2} e^\dagger$  and  $W = \beta_{W,1} e + \beta_{W,2} e^\dagger$  can be written in the idempotent basis as  $Z.W = (\beta_{Z,1} e + \beta_{Z,2} e^\dagger) \cdot (\beta_{W,1} e + \beta_{W,2} e^\dagger) = \beta_{Z,1} \cdot \beta_{W,1} e + \beta_{Z,2} \cdot \beta_{W,2} e^\dagger$ . Thus we see that  $Z$  is invertible if and only if  $\beta_{Z,1} \neq 0 \neq \beta_{Z,2}$  and  $Z^{-1} =$

$\beta_{Z,1}^{-1}e + \beta_{Z,2}^{-1}e^\dagger$ . A non zero  $Z$  that does not have an inverse has the property that either  $\beta_{Z,1} = 0$  or  $\beta_{Z,2} = 0$  and such a  $Z$  is divisor of zero. The zero divisors make up the so called null cone  $NC$  which is closed subset of  $\mathbb{BC}$ . Therefore the set of invertible elements form an open subset of  $\mathbb{BC}$  and every zero divisor is the limit point of the set of regular (invertible) elements in  $\mathbb{BC}$ . The existence of zero divisors in  $\mathbb{BC}$  and the solution of polynomial equation emphasizes the following difference. It comes out that the equation  $\sum_{k=1}^n (a_k + jb_k)(Z = z_1 + z_2j)^k = 0, a_n^2 + b_n^2 \neq 0$  has  $n^2$  solutions in  $\mathbb{BC}$  whereas such type of equation in  $\mathbb{C}$  has  $n$  solutions.

In [8], Hahn Banach Theorem, Closed Graph Theorem, Open Mapping Theorem and Uniform Boundedness Principle for bicomplex Banach spaces were established. For various properties of finite and infinite dimensional bicomplex Hilbert spaces and their applications one can refer to [6], [15] and [16]. Recently D. Alpay, M. E. Luna - Elizarraras, M. Shapiro and D. C. Struppa have written a clear and nice paper on bicomplex function analysis, see [1].

## 2 Bicomplex function spaces and linear operators

In this section we define bicomplex valued function spaces and give some examples of linear operators.

Let  $\Omega = (\Omega, \Sigma, \mu)$  be a  $\sigma$ -finite complete measure space. If  $f = f_1e + f_2e^\dagger$ , where  $f_1$  and  $f_2$  are complex ( $\mathbb{C}(i)$ ) valued measurable functions on  $\Omega = (\Omega, \Sigma, \mu)$ , then  $f$  is a bicomplex valued measurable function on  $\Omega$ . Therefore given any complex ( $\mathbb{C}(i)$ ) valued function space  $(F(\Omega), \|\cdot\|_\Omega)$  we can always define a bicomplex version  $(F(\Omega, \mathbb{BC}), \|\cdot\|_{\mathbb{BC}})$  comprising of all functions of the type  $f = f_1e + f_2e^\dagger$ , where  $f_1$  and  $f_2$  are in  $(F(\Omega), \|\cdot\|_\Omega)$  and  $\|f\|_{\mathbb{BC}} = \frac{1}{\sqrt{2}}(\|f_1\|_\Omega^2 + \|f_2\|_\Omega^2)^{\frac{1}{2}}$ . The addition and scalar multiplication is defined on  $(F(\Omega, \mathbb{BC}), \|\cdot\|_{\mathbb{BC}})$  as under:

$$\begin{aligned} f + g &= (f_1e + f_2e^\dagger) + (g_1e + g_2e^\dagger) \\ &= (f_1 + g_1)e + (f_2 + g_2)e^\dagger \end{aligned}$$

and

$$\begin{aligned} a.f &= (a_1e + a_2e^\dagger).(f_1e + f_2e^\dagger) \\ &= (a_1.f_1)e + (a_2.f_2)e^\dagger, \end{aligned}$$

where  $f, g \in F(\Omega, \mathbb{BC})$  and  $a \in \mathbb{BC}$ . One can easily see that the bicomplex version of complex valued function space is complete if and only if the complex valued function space itself is complete.

**Example 2.1.** Suppose  $L^0(\Omega)$  denotes the linear space of all equivalence classes of complex valued  $\Sigma$ -measurable functions on  $\Omega$  and here any two functions that are equal  $\mu$ -almost everywhere on  $\Omega$  are identified. Then the corresponding bicomplex measurable function space  $L^0(\mathbb{BC})$  comprises of all functions of the type  $f = f_1e + f_2e^\dagger$  where  $f_1, f_2 \in L^0(\Omega)$ .

**Example 2.2.** Suppose  $L^\infty(\Omega)$  denotes the linear space of all equivalence classes of complex valued  $\Sigma$ -measurable essentially bounded functions on  $\Omega$  and here any two functions that are equal  $\mu$ -almost everywhere on  $\Omega$  are identified. Then the corresponding bicomplex measurable function space  $L^\infty(\mathbb{BC})$  comprises of all functions of the type  $f = f_1e + f_2e^\dagger$  where  $f_1, f_2 \in L^\infty(\Omega)$ . Also  $\|f\|_{\infty, \mathbb{BC}} = \|f_1e + f_2e^\dagger\|_{\infty, \mathbb{BC}} = \frac{1}{\sqrt{2}}(\|f_1\|_\infty^2 + \|f_2\|_\infty^2)^{\frac{1}{2}} = \frac{1}{\sqrt{2}}((\text{esssup} f_1)^2 + (\text{esssup} f_2)^2)^{\frac{1}{2}}$

Similarly we can define  $L^p(\mathbb{BC})$ .

**Lemma 2.3.**  $(F(\Omega, \mathbb{BC}), \|\cdot\|_{\mathbb{BC}})$  is complete if and only if  $(F(\Omega), \|\cdot\|_\Omega)$  is complete.

**Proof.** Suppose  $(F(\Omega, \mathbb{BC}), \|\cdot\|_{\mathbb{BC}})$  is complete and  $\{f_n\}$  is a Cauchy sequence in  $(F(\Omega), \|\cdot\|_\Omega)$ . Therefore for given  $\epsilon > 0$  there exists  $r \in \mathbb{N}$  such that

$$\|f_n - f_m\|_\Omega < \epsilon \quad (2.1)$$

for all  $n, m > r$ . Set  $f'_n = f_n e + 0e^\dagger \in \mathbb{BC}$ . Then

$$\begin{aligned} \|f'_n - f'_m\|_{\mathbb{BC}}^2 &= \|(f_n e + 0e^\dagger) - (f_m e + 0e^\dagger)\|_{\mathbb{BC}}^2 \\ &= \|(f_n - f_m)e + 0e^\dagger\|_{\mathbb{BC}}^2 \\ &= \frac{1}{2}(\|f_n - f_m\|_\Omega^2). \end{aligned}$$

This together with (2.1) implies that  $\{f'_n\}$  is a Cauchy in  $(F(\Omega, \mathbb{BC}), \|\cdot\|_{\mathbb{BC}})$ . Therefore by completeness of  $(F(\Omega, \mathbb{BC}), \|\cdot\|_{\mathbb{BC}})$ , there exists  $g = g_1e + g_2e^\dagger$  in  $F(\Omega, \mathbb{BC})$  such that  $f'_n \rightarrow g$  as  $n \rightarrow \infty$ . In order to complete the proof of direct part we need to show that  $f_n \rightarrow g_1$  as  $n \rightarrow \infty$  and  $g_2 = 0$ . Now  $f'_n \rightarrow g$  in  $\mathbb{BC}$ , therefore there exist a natural number  $k$  such that

$$\|f'_n - g\|_{\mathbb{BC}} < \epsilon \quad (2.2)$$

for all  $n > k$ . Also one has

$$\begin{aligned} \|f'_n - g\|_{\mathbb{BC}}^2 &= \|(f_n e + 0e^\dagger) - (g_1e + g_2e^\dagger)\|_{\mathbb{BC}}^2 \\ &= \|(f_n - g_1)e + (0 - g_2)e^\dagger\|_{\mathbb{BC}}^2 \\ &= \frac{1}{2}(\|f_n - g_1\|_\Omega^2 + \|0 - g_2\|_\Omega^2). \end{aligned}$$

This together with (2.2) implies the direct part. For converse, suppose  $(F(\Omega), \|\cdot\|_\Omega)$  is complete and  $\{f_n = f_{n,1}e + f_{n,2}e^\dagger\}_{n=1}^\infty$  is a Cauchy sequence in  $(F(\Omega, \mathbb{BC}), \|\cdot\|_{\mathbb{BC}})$ . Therefore for  $\epsilon > 0$  there exists  $r \in \mathbb{N}$  such that

$$\|f_m - f_n\|_{\mathbb{BC}} < \epsilon \quad (2.3)$$

for all  $m, n > r$ . But

$$\begin{aligned}\|f_m - f_n\|_{\mathbb{B}\mathbb{C}}^2 &= \|(f_{m,1}e + f_{m,2}e^\dagger) - (f_{n,1}e + f_{n,2}e^\dagger)\|_{\mathbb{B}\mathbb{C}}^2 \\ &= \|(f_{m,1} - f_{n,1})e + (f_{m,2} - f_{n,2})e^\dagger\|_{\mathbb{B}\mathbb{C}}^2 \\ &= \frac{1}{2}(\|f_{m,1} - f_{n,1}\|_\Omega^2 + \|f_{m,2} - f_{n,2}\|_\Omega^2)\end{aligned}$$

Therefore by (2.3) we have  $\|f_{m,i} - f_{n,i}\|_\Omega < \sqrt{2}\epsilon$  for  $i = 1, 2$  and for all  $m, n > r$ . This implies that  $\{f_{n,i}\}$  is a Cauchy sequence in  $(F(\Omega), \|\cdot\|_\Omega)$ . Since  $(F(\Omega), \|\cdot\|_\Omega)$  is complete, therefore there exists  $f_i$  in  $(F(\Omega), \|\cdot\|_\Omega)$  such that  $f_{n,i} \rightarrow f_i$  as  $n \rightarrow \infty$  for  $i = 1, 2$ . Therefore there exist natural numbers  $k_1 > 0$  and  $k_2 > 0$  such that

$$\|f_{n,i} - f_i\|_\Omega < \epsilon \quad (2.4)$$

for all  $n > k_i$ . Next we claim that  $f_n = f_{n,1}e + f_{n,2}e^\dagger \rightarrow f = f_1e + f_2e^\dagger$  as  $n \rightarrow \infty$ . Now

$$\begin{aligned}\|f_n - f\|_{\mathbb{B}\mathbb{C}}^2 &= \|(f_{n,1}e + f_{n,2}e^\dagger) - (f_1e + f_2e^\dagger)\|_{\mathbb{B}\mathbb{C}}^2 \\ &= \|(f_{n,1} - f_1)e + (f_{n,2} - f_2)e^\dagger\|_{\mathbb{B}\mathbb{C}}^2 \\ &= \frac{1}{2}(\|f_{n,1} - f_1\|_\Omega^2 + \|f_{n,2} - f_2\|_\Omega^2) \\ &< \epsilon^2\end{aligned}$$

for all  $n > k = \max\{k_1, k_2\}$ . Therefore one has  $\|f_n - f\|_{\mathbb{B}\mathbb{C}} < \epsilon$  for all  $n > k$ . Therefore  $f_n$  is convergent in  $(F(\Omega, \mathbb{B}\mathbb{C}), \|\cdot\|_{\mathbb{B}\mathbb{C}})$ . ■

**Definition 2.4.** A bicomplex linear operator  $T$  is a mapping from a bicomplex normed linear space  $X$  to  $X$  such that  $T(\alpha z + \beta w) = \alpha T(z) + \beta T(w)$ ,  $\forall \alpha, \beta \in \mathbb{B}\mathbb{C}$  and  $z, w \in X$ .

**Remark 2.5.** A bicomplex linear operator  $T : X \mapsto X$  can always be written as  $T = eT_1 + e^\dagger T_2 = e\pi_{i,1}(T) + e^\dagger\pi_{i,2}(T)$ , where  $\pi_{i,l}(T)$  ( $l = 1, 2$ ) is defined as  $\pi_{i,l}(T) = T_l$ .

**Proposition 2.6.** A linear operator  $T : X \mapsto X$  is invertible if and only if both  $T_1$  and  $T_2$  are invertible and  $T^{-1} = eT_1^{-1} + e^\dagger T_2^{-1}$ .

**Proof.** For proof see [3] ■

**Remark 2.7.** Let  $T = T_1e + T_2e^\dagger$  be a bicomplex linear operator on  $(F(\Omega, \mathbb{B}\mathbb{C}), \|\cdot\|_{\mathbb{B}\mathbb{C}})$ .

- (a) A bicomplex linear operator  $T$  is bounded on a bicomplex function space  $(F(\Omega, \mathbb{B}\mathbb{C}), \|\cdot\|_{\mathbb{B}\mathbb{C}})$  if and only if both  $T_1$  and  $T_2$  are bounded on the underlying complex valued function space  $(F(\Omega), \|\cdot\|_\Omega)$ , see [8].
- (b) A bicomplex linear operator  $T$  is compact on a bicomplex function space  $(F(\Omega, \mathbb{B}\mathbb{C}), \|\cdot\|_{\mathbb{B}\mathbb{C}})$  if and only if both  $T_1$  and  $T_2$  are compact on the underlying complex valued function space  $(F(\Omega), \|\cdot\|_\Omega)$ , see [2].

Next we define a bicomplex valued Orlicz space. An Orlicz function  $\phi : [0, \infty) \mapsto [0, \infty]$  is a convex function with  $\phi(0) = 0$  and  $\phi(u) \mapsto \infty$  as  $u \mapsto \infty$  such that  $\phi(u) < \infty$  for some  $0 < u < \infty$ . An Orlicz space  $L^\phi(\Omega)$  is defined as the space of all  $f \in L^0(\Omega)$  such that  $I_\phi(|\lambda f|)$  is finite for some  $\lambda > 0$  and for any  $f \in L^\phi(\Omega)$ , the Orlicz norm of  $f$ ,  $\|f\|_\phi$  is defined as the infimum of all  $\lambda > 0$  such that  $I_\phi(\frac{f}{\lambda}) \leq 1$  where,  $I_\phi(f) = \int_\Omega \phi(|f|) d\mu$ . The Orlicz space  $L^\phi(\Omega)$  is a Banach space with Orlicz norm  $\|\cdot\|_\phi$ .

The bicomplex Orlicz space  $L^\phi(\mathbb{BC})$  is defined as

$$L^\phi(\mathbb{BC}) = \{f_1e + f_2e^\dagger : f_1, f_2 \in L^\phi(\mu)\}.$$

Also the fact that the norm  $\|\cdot\|_{\phi, \mathbb{BC}} : L^\phi(\mathbb{BC}) \mapsto \mathbb{R}$  defined as

$$\|f\|_{\phi, \mathbb{BC}} = \frac{1}{\sqrt{2}}(\|f_1\|_\phi^2 + \|f_2\|_\phi^2)^{\frac{1}{2}}$$

is a complete norm on  $L^\phi(\mathbb{BC})$  follows easily from Lemma 2.3.

**Remark 2.8.** If  $\phi(x) = |x|^p$  for  $p > 1$ , we obtain the bicomplex version of the Lebesgue spaces  $L^p$ .

Now we give some examples of linear operators:

**Definition 2.9.** Let  $T : \Omega \mapsto \Omega$  be a measurable transformation, that is,  $T^{-1}(A) \in \Sigma$  for any  $A \in \Sigma$ . If  $\mu \circ T^{-1}(A) = 0$  for each  $A \in \Sigma$  with  $\mu(A) = 0$ , then  $T$  is said to be non-singular.

Any non-singular measurable transformation  $T$  induces a linear operator  $C_T$  from  $L^0(\Omega)$  into itself defined by

$$(C_T f)(t) = f \circ T(t) = f(T(t)), \quad t \in \Omega, \quad f \in L^0(\Omega).$$

Next if  $f = f_1e + f_2e^\dagger \in L^0(\mathbb{BC})$ , then one can define

$$(C_T f)(t) = f_1 \circ T(t)e + f_2 \circ T(t)e^\dagger$$

**Definition 2.10.** Let  $\Omega = (\Omega, \Sigma, \mu)$  be a  $\sigma$ -finite complete measure space and let  $f = f_1e + f_2e^\dagger \in L^\phi(\mathbb{BC})$ , where  $f_1$  and  $f_2$  are in  $L^\phi$ . For a nonsingular measurable transformation  $T : \Omega \mapsto \Omega$  define composition transformation  $C_T f = C_T(f_1e + f_2e^\dagger) = f_1 \circ T e + f_2 \circ T e^\dagger$ . We say  $C_T$  is composition operator if it maps  $L^\phi(\mathbb{BC})$  into itself.

**Remark 2.11.** Clearly one has  $C_T(f_1e + f_2e^\dagger) = C_T f_1e + C_T f_2e^\dagger$ .

**Remark 2.12.** In light of Remark 2.7, [4] and [17], it easily follows that  $C_T : L^p(\mathbb{BC}) \mapsto L^p(\mathbb{BC})$  is bounded if and only if  $C_T : L^p(\Omega) \mapsto L^p(\Omega)$  is bounded i.e., if and only if

$$\mu \circ T^{-1}(A) \leq M\mu(A),$$

for some  $M > 0$  and for all  $A \in \Sigma$  with  $\mu(A) < \infty$ .

**Example 2.13.** Let  $T : \mathbb{N} \mapsto \mathbb{N}$  be defined by  $T(n) = n - 1$ . Then  $C_T : l^2(\mathbb{BC}) \mapsto l^2(\mathbb{BC})$  is a composition operator on  $l^2(\mathbb{BC})$  defined by  $C_T(x_1, x_2, \dots) = (0, x_1, x_2, \dots)$  and is called the right shift operator.

**Definition 2.14.** Let  $\theta \in L^0(\mathbb{BC})$ . Then we define a multiplication operator  $M_\theta : L^0(\mathbb{BC}) \mapsto L^0(\mathbb{BC})$  as

$$M_\theta f(t) = (\theta(t))(f(t)),$$

for all  $t \in \Omega$  and  $f \in L^0(\mathbb{BC})$ .

**Remark 2.15.** Clearly  $\theta, f \in L^0(\mathbb{BC})$  implies that  $\theta = \theta_1 e + \theta_2 e^\dagger$  for some  $\theta_1, \theta_2 \in L^0(\Omega)$  and  $f = f_1 e + f_2 e^\dagger$  for some  $f_1, f_2 \in L^0(\Omega)$ . Therefore

$$\begin{aligned} M_\theta &= (\theta)(f) \\ &= (\theta_1 e + \theta_2 e^\dagger)(f_1 e + f_2 e^\dagger) \\ &= (\theta_1)(f_1)e + (\theta_2)(f_2 e^\dagger) \\ &= (M_{\theta_1} f_1)e + (M_{\theta_2} f_2)e^\dagger \end{aligned}$$

where  $M_{\theta_1}$  and  $M_{\theta_2}$  are multiplication operators on  $L^0(\Omega)$ . Therefore  $M_\theta$  is bounded on  $L^0(\mathbb{BC})$  if and only if  $M_{\theta_1}$  and  $M_{\theta_2}$  are bounded on  $L^0(\Omega)$  i.e., if and only if  $\theta_1, \theta_2 \in L^\infty(\Omega)$ .

**Lemma 2.16.** Let  $1 < p < \infty$ . Then the sequence of unit vectors  $\{e_1, e_2, e_3, \dots\}$ , where  $e_k = \delta_{kj}$  is a Schauder basis for  $l^p(\mathbb{BC})$ .

**Proof.** For any  $x = \{x_n\} \in l^p(\mathbb{BC})$ ,

$$\begin{aligned} \|x - \sum_{k=1}^n x_k e_k\| &= \|\{x_{n+1}, x_{n+2}, x_{n+3}, \dots\}\| \\ &= \left( \sum_{k=n+1}^{\infty} |x_k|^p \right)^{\frac{1}{p}} \\ &= \left( \sum_{k=n+1}^{\infty} |e x_{k1} + e^\dagger x_{k2}|^p \right)^{\frac{1}{p}} \\ &\leq \left( \sum_{k=n+1}^{\infty} |e x_{k1}|^p + \sum_{k=n+1}^{\infty} |e^\dagger x_{k2}|^p \right)^{\frac{1}{p}} \\ &= \left( \sum_{k=n+1}^{\infty} |e|^p |x_{k1}|^p + \sum_{k=n+1}^{\infty} |e^\dagger|^p |x_{k2}|^p \right)^{\frac{1}{p}} \\ &\rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

Therefore  $x = \sum_{k=1}^{\infty} x_k e_k$  and hence  $\{e_1, e_2, e_3, \dots\}$  is a Schauder basis for  $l^p(\mathbb{BC})$ . ■

**Theorem 2.17.** The dual space of  $l^p(\mathbb{BC})$  is  $l^q(\mathbb{BC})$  where  $1 < p < \infty$  and  $\frac{1}{p} + \frac{1}{q} = 1$ .

**Proof.** Since by Lemma 2.16, the sequence of unit vectors  $\{e_1, e_2, e_3, \dots\}$ , where  $e_k = \delta_{kj}$  is a Schauder basis for  $l^p(\mathbb{BC})$ . The rest of the proof follows on the similar lines as in the case of  $l^p(\Omega)$ . ■

Next we define modular  $I_{\phi, \mathbb{BC}}$  on bicomplex Orlicz space as

$$I_{\phi, \mathbb{BC}}(f) = I_\phi(f_1)e + I_\phi(f_2)e^\dagger \tag{2.5}$$

**Theorem 2.18.** Let  $f = f_1e + f_2e^\dagger \in L^\phi(\mathbb{BC})$ , where  $f_1$  and  $f_2$  are in  $L^\phi(\mu)$ .

(a) If  $\|f\|_{\phi, \mathbb{BC}} \leq \frac{1}{\sqrt{2}}$  then  $I_\phi(f_i) \leq 1$ , for  $i = 1, 2$ .

(b) If  $I_\phi(f_i) \leq 1$ , for  $i = 1, 2$  then  $\|f\|_{\phi, \mathbb{BC}} \leq 1$ .

**Proof.**

(a) Let  $\|f\|_{\phi, \mathbb{BC}} = \frac{1}{\sqrt{2}}(\|f_1\|_\phi^2 + \|f_2\|_\phi^2)^{\frac{1}{2}} \leq \frac{1}{\sqrt{2}}$ . This implies that  $\|f\|_{\phi, \mathbb{BC}}^2 = \frac{1}{2}(\|f_1\|_\phi^2 + \|f_2\|_\phi^2) \leq \frac{1}{2}$ . Which yields that  $\|f_i\|_\phi^2 \leq 1$ , for  $i = 1, 2$ . Therefore  $I_\phi(f_i) \leq 1$ , for  $i = 1, 2$ .

(b) Let  $I_\phi(f_i) \leq 1$ , for  $i = 1, 2$ . This yields that  $\|f_i\|_\phi \leq 1$ , for  $i = 1, 2$ . Therefore  $\|f\|_{\phi, \mathbb{BC}} = \frac{1}{\sqrt{2}}(\|f_1\|_\phi^2 + \|f_2\|_\phi^2)^{\frac{1}{2}} \leq 1$ .  $\blacksquare$

A particular case of  $L^\phi(\mathbb{BC})$  is  $l^\phi(\mathbb{BC})$  which is defined as

$$l^\phi(\mathbb{BC}) = \{\{x_n\}_{n=1}^\infty : x_n = x_{n,1}e + x_{n,2}e^\dagger \in \mathbb{BC} \text{ and } \{x_{n,i}\}_{n=1}^\infty \in l^\phi(\mathbb{C}) \text{ for } i = 1, 2.\}$$

and  $\|\{x_n\}\|_{\phi, \mathbb{BC}} = \frac{1}{\sqrt{2}}(\|\{x_{n,1}\}\|_\phi^2 + \|\{x_{n,2}\}\|_\phi^2)^{\frac{1}{2}}$ . The addition and scalar multiplication is defined component wise on  $l^\phi(\mathbb{BC})$ .

**Theorem 2.19.**  $l^\phi(\mathbb{BC})$  equipped with  $\|\cdot\|_{\phi, \mathbb{BC}}$  is a Banach space over  $\mathbb{BC}$ .

**Proof.** It is clear from the definition of addition and scalar multiplication that  $l^\phi(\mathbb{BC})$  is a module over  $\mathbb{BC}$ . It is also easy to show that  $\|\cdot\|_{\phi, \mathbb{BC}}$  defines a norm on  $l^\phi(\mathbb{BC})$ . Now it only remains to show that  $l^\phi(\mathbb{BC})$  is also complete with respect  $\|\cdot\|_{\phi, \mathbb{BC}}$ . For that let  $\{x_n^k\}_{k=1}^\infty$  be a Cauchy sequence in  $l^\phi(\mathbb{BC})$ . Since for  $\{x_n = x_{n,1}e + x_{n,2}e^\dagger\}_{n=1}^\infty$  we have  $\|\{x_n\}\|_{\phi, \mathbb{BC}} = \frac{1}{\sqrt{2}}(\|\{x_{n,1}\}\|_\phi^2 + \|\{x_{n,2}\}\|_\phi^2)^{\frac{1}{2}}$ . Therefore one has

$$\|\{x_{n,i}\}\|_\phi \leq \sqrt{2}\|\{x_n\}\|_{\phi, \mathbb{BC}} \text{ for } i = 1, 2. \quad (2.6)$$

Now for  $\epsilon > 0$  there exists  $r \in \mathbb{N}$  Such that

$$\|\{x_n^l - x_n^m\}\|_{\phi, \mathbb{BC}} < \frac{\epsilon}{\sqrt{2}} \text{ for all } l, m \geq r. \quad (2.7)$$

Therefore (2.6) and (2.7) yield that  $\{x_{n,i}^k\}_{k=1}^\infty$  is a Cauchy sequence in  $(l^\phi(\mathbb{C}), \|\cdot\|_\phi)$  for  $i = 1, 2$ . In view of the completeness of the  $(l^\phi(\mathbb{C}), \|\cdot\|_\phi)$ , we have that  $\{x_{n,i}^k\}_{k=1}^\infty$  converges to some  $\{x_{n,i}\} \in l^\phi(\mathbb{C})$ . Therefore for  $i = 1, 2$  there exists  $p_i \in \mathbb{N}$  such that

$$\|\{x_{n,i}^l - x_{n,i}\}\|_\phi < \sqrt{\epsilon} \text{ for all } l \geq p_i$$

Now for  $p = \max\{p_1, p_2\}$  and  $l \geq p$  one has

$$\begin{aligned} \|\{x_n^l - x_n\}\|_{\phi, \mathbb{BC}}^2 &= \frac{1}{2}(\|\{x_{n,i}^l - x_{n,i}\}\|_\phi^2 + \|\{x_{n,i}^l - x_{n,i}\}\|_\phi^2) \\ &< \frac{1}{2}(\epsilon + \epsilon) \\ &= \epsilon. \end{aligned}$$

Thereby showing that  $\{x_n^k\}_{k=1}^\infty$  is a convergent sequence in  $l^\phi(\mathbb{BC})$ . Therefore  $l^\phi(\mathbb{BC})$  equipped with  $\|\cdot\|_{\phi, \mathbb{BC}}$  is a Banach space over  $\mathbb{C}$ .  $\blacksquare$



**Theorem 2.20.** Let  $(\Omega, \Sigma, \mu)$  be a  $\sigma$ -finite and purely atomic measure space with atoms  $\{A_n\}$  with measure  $\mu(A_n) = a_n > 0$  for any  $n \in \mathbb{N}$ . Let  $T : \Omega \mapsto \Omega$  be a measurable transformation with  $T(\Omega) = \Omega$  and  $b_n = \frac{\mu \circ T^{-1}(A_n)}{\mu(A_n)}$ . Then  $C_T$  is bounded from  $l^\phi(\mathbb{BC}, \{a_n\})$  into itself if and only if for any  $\{x_n\}$  in  $l^\phi(\mathbb{BC}, \{a_n\})$ , there exist scalars  $\lambda_1$  and  $\lambda_2$  such that

$$\sum_{n=1}^{\infty} (\phi(\lambda_1 x_{n,1}) + \phi(\lambda_2 x_{n,2})) b_n < \infty \quad (2.8)$$

where  $x_n = x_{n,1}e + x_{n,2}e^\dagger$ .

**Proof.** Suppose  $C_T : l^\phi(\mathbb{BC}, \{a_n\}) \mapsto l^\phi(\mathbb{BC}, \{a_n\})$  is bounded. Therefore there exists a  $k > 0$  such that

$$\|C_T(\{x_n\})\|_{\phi, \mathbb{BC}} \leq k \|\{x_n\}\|_{\phi, \mathbb{BC}} \quad (2.9)$$

for every  $\{x_n\} \in l^\phi(\mathbb{BC}, \{a_n\})$ . Now set  $x_n^1$  and  $x_n^2$  for  $x_n = x_{n,1}e + x_{n,2}e^\dagger \in \mathbb{BC}$  as

$$\begin{aligned} x_n^1 &= x_{n,1}e + 0e^\dagger \\ x_n^2 &= 0e + x_{n,2}e^\dagger. \end{aligned}$$

Then clearly  $\{x_n^1\}$  and  $\{x_n^2\}$  belong to  $l^\phi(\mathbb{BC}, \{a_n\})$  whenever  $\{x_n\} \in l^\phi(\mathbb{BC}, \{a_n\})$  and

$$\begin{aligned} x_{n,1}^1 &= x_{n,1}, & x_{n,1}^2 &= 0; \\ x_{n,2}^1 &= 0, & x_{n,2}^2 &= x_{n,2}. \end{aligned}$$

Therefore, by (2.9), if  $\{x_n\} \in l^\phi(\mathbb{BC}, \{a_n\})$  we have

$$\|C_T(\{x_n^i\})\|_{\phi, \mathbb{BC}} \leq \|\{x_n^i\}\|_{\phi, \mathbb{BC}}$$

for  $i = 1, 2$ . Therefore,  $\|\{x_{T(n),1}^i\}\|_\phi \leq \sqrt{k} \|\{x_{n,1}^i\}\|_\phi$ . Consequently we have  $C_T : l^\phi(\{a_n\}) \mapsto l^\phi(\{a_n\})$  is bounded. Therefore, by Theorem 2.6 in [4], we have  $\lambda_i > 0$  such that

$$\sum_{n=1}^{\infty} (\lambda_i x_{n,i}) b_n < \infty \quad (2.10)$$

for  $i = 1, 2$ . Now (2.10) easily yields (2.8).

Conversely suppose that (2.8) holds. Then there exists  $\lambda_i > 0$  such that

$$\sum_{n=1}^{\infty} (\lambda_i x_{n,i}) b_n < \infty \quad (2.11)$$

for  $i = 1, 2$  and  $\{x_n = x_{n,1}e + x_{n,2}e^\dagger\} \in l^\phi(\mathbb{BC}, \{a_n\})$ . Again by Theorem 2.6 in [4],  $C_T : l^\phi(\{a_n\}) \mapsto l^\phi(\{a_n\})$  is bounded. Therefore there exists  $k_i > 0$  such that

$$\|\{x_{T(n),i}\}\|_\phi \leq k_i \|\{x_{n,i}\}\|_\phi.$$

Therefore one has

$$\begin{aligned}
\|C_T(\{x_n\})\|_{\phi, \mathbb{BC}}^2 &= \|\{x_{T(n)}\}\|_{\phi, \mathbb{BC}}^2 \\
&= \frac{1}{2}(\|\{x_{T(n),1}\}\|_{\phi}^2 + \|\{x_{T(n),2}\}\|_{\phi}^2) \\
&\leq \frac{k_1}{2}\|\{x_{n,1}\}\|_{\phi}^2 + \frac{k_2}{2}\|\{x_{n,2}\}\|_{\phi}^2 \\
&= \frac{k}{2}(\|\{x_{n,1}\}\|_{\phi}^2 + \|\{x_{n,2}\}\|_{\phi}^2) \\
&= k\|\{x_n\}\|_{\phi, \mathbb{BC}}^2
\end{aligned}$$

where  $k = \max \{k_1, k_2\}$ . Therefore  $C_T$  is bounded from  $l^\phi(\mathbb{BC}, \{a_n\})$  into itself. ■

**Definition 2.21.** An Orlicz function  $\phi$  is called an N - function if

1.  $\phi$  is continuous.
2.  $\phi(x) = 0$  if and only if  $x = 0$ .
3.  $\lim_{x \rightarrow 0} \frac{\phi(x)}{x} = 0$ .
4.  $\lim_{x \rightarrow \infty} \frac{\phi(x)}{x} = \infty$ .

**Theorem 2.22.** If  $\tilde{L}^\phi(\mathbb{BC}) = \{f_1e + f_2e^\dagger : f_1, f_2 \in \tilde{L}^\phi\}$  then

1.  $\bigcup \{\tilde{L}^\phi(\mathbb{BC}) : \phi \text{ ranges over all N - functions}\} \subseteq L^1(\mathbb{BC})$ .
2.  $L^\infty(\mathbb{BC}) = \bigcap \{\tilde{L}^\phi(\mathbb{BC}) : \phi \text{ ranges over all N - functions}\}$ .

**Proof.**

1. If  $f = f_1e + f_2e^\dagger \in \bigcup \{\tilde{L}^\phi(\mathbb{BC}) : \phi \text{ ranges over all N - functions}\}$ , then  $f = f_1e + f_2e^\dagger \in \tilde{L}^\phi(\mathbb{BC})$  for some  $\phi$ . Therefore  $f_1$  and  $f_2$  are in  $\tilde{L}^\phi$  for some  $\phi$ . This implies that  $f_1$  and  $f_2$  are in  $L^1(\mu)$ . Hence  $f = f_1e + f_2e^\dagger \in L^1(\mathbb{BC})$ . Therefore  $\bigcup \{\tilde{L}^\phi(\mathbb{BC}) : \phi \text{ ranges over all N - functions}\} \subseteq L^1(\mathbb{BC})$ .
2. Let  $f = f_1e + f_2e^\dagger \in L^\infty(\mathbb{BC})$ , where  $f_1$  and  $f_2$  are in  $L^\infty(\mu)$ . Therefore  $f_1$  and  $f_2$  are in  $\tilde{L}^\phi$  for all  $\phi$ . This implies that  $f = f_1e + f_2e^\dagger \in \tilde{L}^\phi(\mathbb{BC})$  for all  $\phi$ . Hence  $f = f_1e + f_2e^\dagger \in \bigcap \{\tilde{L}^\phi(\mathbb{BC}) : \phi \text{ ranges over all N - functions}\}$ . Conversely if  $f = f_1e + f_2e^\dagger \in \bigcap \{\tilde{L}^\phi(\mathbb{BC}) : \phi \text{ ranges over all N - functions}\}$ , then  $f = f_1e + f_2e^\dagger \in \tilde{L}^\phi(\mathbb{BC})$  for all  $\phi$ . Therefore  $f_1$  and  $f_2$  are in  $\tilde{L}^\phi$  for all  $\phi$ . This implies that  $f_1$  and  $f_2$  are in  $L^\infty(\mu)$ . Hence  $f = f_1e + f_2e^\dagger \in L^\infty(\mathbb{BC})$ . Therefore  $L^\infty(\mathbb{BC}) = \bigcap \{\tilde{L}^\phi(\mathbb{BC}) : \phi \text{ ranges over all N - functions}\}$ . ■

**Theorem 2.23.** Let  $f_n = f_{n,1}e + f_{n,2}e^\dagger, n \geq 1$  be a sequence in  $L^\phi(\mathbb{BC})$  such that  $f_n \rightarrow f = f_1e + f_2e^\dagger$ . Then  $\|f\|_{\mathbb{BC}} \leq \varliminf_{n \rightarrow \infty} \|f_n\|_{\mathbb{BC}}$ .

**Proof.** The result follows immediately from

$$\begin{aligned}
2\|f\|_{\mathbb{B}\mathbb{C}}^2 &= \|f_1\|_{\Omega}^2 + \|f_2\|_{\Omega}^2 \\
&\leq \underline{\lim}_{n \rightarrow \infty} \|f_{n,1}\|_{\Omega}^2 + \underline{\lim}_{n \rightarrow \infty} \|f_{n,2}\|_{\Omega}^2 \\
&\leq \underline{\lim}_{n \rightarrow \infty} (\|f_{n,1}\|_{\Omega}^2 + \|f_{n,2}\|_{\Omega}^2) \\
&= 2 \underline{\lim}_{n \rightarrow \infty} \|f_n\|_{\mathbb{B}\mathbb{C}}^2.
\end{aligned}$$

■

**Theorem 2.24.** *Let  $\Omega = (\Omega, \Sigma, \mu)$  be a  $\sigma$ -finite complete measure space and  $T : \Omega \mapsto \Omega$  be a non singular measurable transformation. Also suppose that  $\phi$  satisfies  $\Delta_2$  condition. Then  $C_T : L^\phi(\mathbb{B}\mathbb{C}) \mapsto L^\phi(\mathbb{B}\mathbb{C})$  is bounded if and only if there is a constant  $m > 0$  such that  $\mu \circ T^{-1}(A) \leq m\mu(A)$  for each  $A$  such that  $\mu(A) < \infty$ .*

**Proof.** First suppose that there is a constant  $m > 0$  such that  $\mu \circ T^{-1}(A) \leq m\mu(A)$  for each  $A$  such that  $\mu(A) < \infty$ . Therefore there exists a constant  $k$  such that

$$\begin{aligned}
2\|C_T f\|_{\mathbb{B}\mathbb{C}}^2 &= 2\|C_T f_1 e + C_T f_2 e^\dagger\|_{\mathbb{B}\mathbb{C}}^2 \\
&= \|C_T f_1\|_{\Omega}^2 + \|C_T f_2\|_{\Omega}^2 \\
&\leq k\|f_1\|_{\Omega}^2 + k\|f_2\|_{\Omega}^2 \\
&= k 2 \|f\|_{\mathbb{B}\mathbb{C}}^2,
\end{aligned}$$

Therefore  $\|C_T f\|_{\mathbb{B}\mathbb{C}} \leq \sqrt{k}\|f\|_{\mathbb{B}\mathbb{C}}$ . Hence  $C_T$  is bounded. Conversely suppose that  $C_T$  is bounded on  $L^\phi(\mathbb{B}\mathbb{C})$ . Then it follows from Remark 2.7 that  $C_T$  is bounded on  $L^\phi(\Omega)$  and therefore the result follows from [4]. ■

## Acknowledgment

The first author is thankful to D. Rochon for motivating him to work on bicomplex functional analysis during his several visits to University of Jammu.

## References

- [1] D. Alpay, M. E. Luna - Elizarraras, M. Shapiro, D. C. Struppa, *Basics of functional analysis with bicomplex scalars, and bicomplex Schur analysis*, (2013), arXiv: 1304.0781v1[math.FA].
- [2] K.S. Charak, Ravinder. Kumar and D. Rochon, *Infinite dimensional bicomplex spectral decomposition theorem*, Adv. Appl. Clifford Algebras, SP Birkhauser Verlag Basel, DOI 10.1007/s00006-013-0385-5 (2013).
- [3] F. Colombo, I. Sabadini, D.C. Struppa, *Bicomplex holomorphic functional calculus*, Math. Nachr. 1 – 13 (2013), DOI 10.1002/mana.201200354.
- [4] Y. Cui, H. Hudzik, R. Kumar and L. Maligranda, *Composition operators in Orlicz spaces*, J. Austral. Math. Soc., **76** (2004), 189–206.

- [5] R. Gervais Lavoie, L. Marchildon and D. Rochon, *Infinite-Dimensional Bicomplex Hilbert Spaces*, Ann. Funct. Anal, **1** No. 2 (2010), 75–91.
- [6] R. Gervais Lavoie and D. Rochon, *The Bicomplex Heisenberg Uncertainty Principle, Theoretical Concepts of Quantum Mechanics*, InTech Book, (2012), 39–64.
- [7] R. Gervais Lavoie, L. Marchildon and D. Rochon, *Hilbert Space of the Bicomplex Quantum Harmonic Oscillator*, AIP Conference Proceedings **1327**(2011), 148-157 .
- [8] Romesh. Kumar, Rajeev. Kumar and D. Rochon, *The fundamental theorems in the framework of Bicomplex Topological Modules*, (2011) arXiv: 1109.3424v1[math.FA].
- [9] J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces I and II*, Springer - Verlag, Berlin Heidelberg, 1996.
- [10] M.E. Luna-Elizarraras, M. Shapiro, D.C. Struppa and A. Vajiac, *Bicomplex numbers and their elementary functions*, Cubo, A Mathematical Journal, **14**No.2 (2013),61–80,.
- [11] M.E. Luna-Elizarraras, M. Shapiro, D.C. Struppa and A. Vajiac, *Complex Laplacian and derivatives of bicomplex functions*, Complex Anal. Oper. Theory, DOI 10.1007/s11785-013-0284-8(2013).
- [12] L. Maligranda, *Orlicz Spaces and Interpolation*, Seminars in Math. **5**, Univ. Estadual de Campinas, Campinas SP, Brazil, 1989.
- [13] G. B. Price, *An introduction to multicomplex spaces and functions*, 3rd edition, Marcel Dekker New York, 1987.
- [14] M. M. Rao and Z. D. Ren, *Theory of Orlicz spaces*, Marcel Dekker, New York, 1991.
- [15] D. Rochon and M. Shapiro, ‘On algebraic properties of bicomplex and hyperbolic numbers’, Anal. Univ. Oradea, fasc. math., **11** (2004), 71–110.
- [16] D. Rochon and S. Tremblay, *Bicomplex Quantum Mechanics II: The Hilbert Space*, Advances in applied Clifford algebras, **16** No. 2 (2006), 135–157.
- [17] R. K. Singh, *Composition operators induced by rational functions*, Proc. Amer. Math. Soc., **59** No. 2(1976), 329–333.

Romesh Kumar, *Department of Mathematics, University of Jammu, Jammu, J&K - 180 006, India.*

E-mail : romesh\_jammu@yahoo.com

Kailash Sharma, *Department of Mathematics, University of Jammu, Jammu, J&K - 180 006, India.*

E-mail :kailashsharma\_ju@yahoo.co.in

Sumit Dubey, *Department of Mathematics, University of Jammu, Jammu, J&K - 180 006, India.*

E-mail : sumitdubey@live.in